

## References

1. Igner J, Loske A. Zur Etymologie des Wortes „Schwindel“. *HNO aktuell* 1997; 12: 339-340
2. Reker U, Frese KA. Objektive Kriterien für die Begutachtung des vestibulären Schwindels. *Laryngo Rhino Otol* 1998; 77: 179-184
3. Probst R. Schwindel aus neurootologischer Sicht. *Therapeutische Umschau* 1995; 52: 724-731
4. Schaaf H, Nelting M, Hesse G. Schwindel-psychosomatisch gesehen. München: Profil, 1999
5. Dieterich M, Eckardt-Henn A. Neurologische und somatoforme Schwindelsyndrome. *Nervenarzt* 2004; 75: 281-302
6. Schaaf H. Psychogener Schwindel in der HNO-Heilkunde. *HNO* 2001; 49: 307-315
7. Langman J. Medizinische Embryologie. Stuttgart: Thieme 1977
8. Murofushi T, Ouvrier RA, Parker GD, Graham RL, da Silva M, Halmagyi GM. Vestibular abnormalities in charge association. *Ann Otol Rhinol Laryngol* 1997; 106: 129-134
9. Lemmerling MM, Vanzieleghem BD, Mortier GR, Dhooge IJ, Kunnen MF. Unilateral semicircular canal aplasia in Goldenhar's syndrome. *Am J Neuroradiol* 2000; 21: 1334-1336
10. Lang J. Anatomie des vestibulären Systems. In: Haid CT (Hrsg). Schwindel aus interdisziplinärer Sicht. Stuttgart: Thieme 2003; 4-16
11. Rennie KJ, Manning KC, Ricci AJ. Mechano-electrical transduction in the turtle utricle. *Biomed Sci Instrum* 2004; 40: 441-44
12. Spoedlin H. Strukturelle Organisation des Innenohres. In: Oto-Rhino-Laryngologie in Klinik und Praxis. Bd 1. (Hrsg. Helms J, Herberhold C, Kasenbauer E). Stuttgart: Thieme; 1994; 32-74
13. Reiss G, Wolkowiak W, Zenner HP, Plinkert PK, Lehnhardt E. Das stato-akustische Organ. Hannover: Duphar Pharma, 1989
14. Scherer H. Das Gleichgewicht. Berlin: Springer, 1997
15. Brandt T, Dieterich M, Strupp M. Vertigo. Darmstadt: Steinkopff, 2004
16. Brandt T, Bartenstein P, Janek A, Dieterich M. Reciprocal inhibitory visual-vestibular interaction. Visual motion stimulation deactivates the parieto-insular vestibular cortex. *Brain* 1998; 121: 1749-1758
17. Dieterich M, Brandt T. Brain activation studies on visual-vestibular and ocular motor interaction. *Curr Opin Neurol* 2000; 13: 13-18
18. Dieterich M, Brandt T. Vestibular system: anatomy and functional magnetic resonance imaging. *Neuroimaging Clin N Am* 2001; 11: 263-273
19. Brandt T, Glasauer S, Stephan T, Bense S, Yousry TA, Deutschlander A, Dieterich M. Visual-vestibular and visuo-visual cortical interaction: new insights from fMRI and pet. *Ann N Y Acad Sci* 2002; 956: 230-241
20. Kleinschmidt A, Thilo KV, Büchel C, Gresty MA, Bronstein AM, Frackowiak RS. Neural correlates of visual-motion perception as object- or self-motion. *Neuroimage*. 2002; 16: 873-882
21. Dieterich M, Bense S, Stephan T, Yousry TA, Brandt T. fMRI signal increases and decreases in cortical areas during small-field optokinetic stimulation and central fixation. *Exp Brain Res* 2003; 148: 117-127
22. Naidich TP, Kang E, Fatterpekar GM, Delman BN, Gultekin SH, Wolfe D, Ortiz O, Yousry I, Weismann M, Yousry TA. The insula: anatomic study and MR imaging display at 1.5 T. *AJR Am J Neuroradiol* 2004; 25: 222-232
23. Brandt T, Dieterich M. Central vestibular syndromes in roll, pitch and yaw planes. Topographic diagnosis of brainstem disorders. *Neuroophthalmology* 1995; 15: 291-303
24. Leblanc A. Atlas of hearing and balance organs. Paris: Springer, 1999
25. Schreiner L. Neue experimentelle und klinische Erkenntnisse zur Frage einer interlabyrinthären Verbindung. *Laryngo Rhino Otol* 1999, 78; 387-393
26. Harris JP, Low NC, House WF. Contralateral hearing loss following inner ear injury: sympathetic cochleolabyrinthitis? *Am J Otol* 1985; 6: 371-377
27. Salt AN, Rask-Andersen H. Responses of the endolymphatic sac to perilymphatic injections and withdrawals: evidence for the presence of a one-way valve. *Hear Res* 2004; 191: 90-100
28. Schuknecht HF, Ruther A. Blockage of longitudinal flow in endolymphatic hydrops. *Eur Arch Otorhinolaryngol* 1991; 248: 209-217
29. Westhofen M. Vestibulärer Schwindel. Neue Entwicklungen zwischen Otologie, Neurologie und Otochirurgie. *Laryngo Rhino Otol* 2001; 80: 651
30. Walther LE, Schmidt WD, Gudziol H, Scheibe A, Scheiding B, Rössler B, Fassler D, Beleites E. Nahinfarotreizung des Gleichgewichtsorgans-erste klinische Erfahrungen. *Laryngo Rhino Otol* 2003; 82: 687-692
31. Walther LE, Gudziol H, Beleites E. Gezielte Strahlungskalorisation des Gleichgewichtsorgans mittels Nahinfarot. *Laryngo Rhino Otol* 2004; 83: 88-95
32. Walther LE, Scheibe A, Schmidt WD, Römhild D, Fassler D, Gudziol H, Beleites E. Untersuchungen zur thermischen Reizung des Gleichgewichtsorgans mittels Wärmestrahlung (NIR). *HNO* 2004; 52: 525-532
33. Halmagyi GM; Curthoys IS: A clinical sign of canal paresis. *Arch Neurol* 1988; 45: 737-739
34. Halmagyi GM, Aw ST, Karlberg M, Curthoys IS, Todd MJ. Inferior vestibular neuritis. *Ann N Y Acad Sci* 2002; 956: 306-313
35. Westhofen M. Aktueller Stand der Diagnostik und der operativen Therapie von Otolithenerkrankungen. In: Stoll W (Hrsg). *Vestibuläre Erkrankungen-eine interdisziplinäre Herausforderung*. Stuttgart: Thieme, 2001, 83-97

36. Kömpf D. Der pseudovestibuläre Kleinhirninsult. Nervenarzt 1986; 57: 163-166
37. Grad A, Baloh RW. Vertigo of vascular origin. Clinical and electronystagmographic features in 84 cases. Arch Neurol 1989; 46: 281-284
38. Gomez CR, Cruz-Flores S, Malkoff MD, Sauer CM, Burch CM. Isolated vertigo as a manifestation of vertebrobasilar ischemia. Neurology. 1996; 47: 94-97.
39. Berlit P. Isolierter Schwindel bei vertebrobasilärer Ischämie-gibt es das? HNO 1998; 46: 296-300
40. Walther LE, Beleites E. Schwindeldiagnostik. Ärztebl Thüring 1999; 10: 549-554
41. Schmiz A, Haibt-Lüttke G, Albrecht G, Hagen R, Gustorf-Aeckerle R. Die Thrombose der Arteria basilaris-eine seltene Differenzialdiagnose der Hörsturzes und des Vestibularisausfalls. Laryngo Rhino Otol 2000; 79: 253-259
42. Westhofen M. Hals-Nasen-Ohrenheilkunde systematisch. Bremen: Uni-Med 2001
43. Whitehead RE, MacDonald CB, Melhem ER, McMahon L. Spontaneous labyrinthine hemorrhage in sickle cell disease. AJNR Am J Neuroradiol. 1998; 19: 1437-1440
44. Stange T. Hämorrhagische Labyrinthitis: Ursache für akuten Innenohrausfall. Vortrag XIX. Neusser HNO-Kolloquium 2004
45. Helling K, Clarke AH, Watanabe N, Scherer H. Morphologische Untersu- chungen zur Form der Cupula in der Bogengangssampulle. HNO 2000; 48: 822-827
46. Scherer H, Helling K, Watanabe S. Zur Pathogenese der sogenannten Neuropathia vestibularis-Experimente an der Ampulle. In: Stoll W (Hrsg). Vestibuläre Erkrankungen-eine interdisziplinäre Herausforderung. Stuttgart: Thieme, 2001, 16-22
47. Helling K, Watanabe N, Jijiwa H, Mizuno Y, Watanabe S, Scherer H. Altered cupular mechanics: a cause of peripheral vestibular disorders? Acta Otolaryngol 2002; 122: 386-391
48. Arbusow V, Theil D, Strupp M, Mascalo A, Brandt T. HSV-1 not only in human vestibular ganglia but also in the vestibular labyrinth. Audiol Neurotol 2001; 6: 259-262
49. Gacek RR, Gacek MR. Vestibular neuritis: a viral neuropathy. In: Arnold W (Hrsg). Viral neuropathies in the temporal bone. Adv Otorhinolaryngol 2002; 60: 54-66
50. Fenton JE, Shirazi A, Turner J, Fagan PA. Atypical vestibular neuritis: a case report. Otolaryngol Head Neck Surg 1995; 112: 738-741
51. Strupp M, Jager L, Müller-Lisse U, Arbusow V, Reiser M, Brandt T. High resolution Gd-DTPA MR imaging of the inner ear in 60 patients with idiopathic vestibular neuritis: no evidence for contrast enhancement of the labyrinth or vestibular nerve. J Vestib Res. 1998; 8: 427-433
52. Shimizu T, Sekitani T, Hirata T, Hara H. Serum viral antibody titer in vestibular neuronitis. Acta Otolaryngol Suppl 1993; 503:74-78
53. Walther LE, Hentschel H, Oehme A, Gudziol H, Beleites E. Herpesviren bei akuter Innenohrschwerhörigkeit und "Vestibularisausfall": klinische und serologische Befunde. Otorhinolaryngol Nova 2002; 12: 124-131
54. Haid T, Mirsberger J. Die periphere Neuropathia vestibularis und ein zentral-vestibuläres Äquivalent. HNO 1985; 33: 262-270
55. Ellies M, Politycki C, Laskawi R. Therapie des isolierten einseitigen Vestibularisausfalls. Otorhinolaryngol Nova 1996; 6: 264-270
56. Herzog N, Allum JJJ, Probst R. Verlauf der kalorischen Erregbarkeit nach akuter peripherer vestibulärer Funktionsstörung. HNO 1997; 45: 123-127
57. Strupp M, Zingler VC, Arbusow V, Niklas D, Maag KP, Dieterich M, Bense S, Theil D, Jahn K, Brandt T. Methylprednisolone, valacyclovir, or the combination for vestibular neuritis. N Engl J Med. 2004; 351: 354-361
58. Gacek RR, Gacek MR. Antiviral therapy of vestibular ganglionitis. In: Arnold W (Hrsg). Viral neuropathies in the temporal bone. Adv Otorhinolaryngol 2002, 60. 124-126
59. Scholtz AW, Schwarz M, Baumann W, Kleinfeldt D, Scholtz HJ. Treatment of vertigo due to acute unilateral vestibular loss with a fixed combination of cinnarizine and dimenhydrinate: a double-blind, randomized, parallel-group clinical study. Clin Ther 2004; 26: 866-877
60. Schneider D, Kiessling B, Wieczorek M, Bognar-Steinberg I, Schneider L, Claussen CF. Influence of 3 antivertiginous medications on the vigilance of healthy volunteers. Int J Clin Pharmacol Ther 2003; 41: 171-181
61. Weiser M, Strosser W, Klein P. Homeopathic vs conventional treatment of vertigo: a randomized double-blind controlled clinical study. Arch Otolaryngol Head Neck Surg. 1998; 124: 879-885
62. Westhofen M. Verlauf der kalorischen Erregbarkeiten nach akuter peripherer Funktionsstörung. HNO 1997; 45: 112-113
63. Igarashi M. Compensation for peripheral vestibular disturbances-animal studies. In Bles W, Brandt T (Hrsg). Disorders of posture and gait. Elsevier: Amsterdam, 1986, 337-351
64. Gramowski KH. Das trainierte Labyrinth. HNO (Berl) 1964; 12: 105
65. Ehrenberger K. Prinzipien einer konservativen Therapie peripherer und zentraler Gleichgewichtsstörungen. HNO 1988; 36: 301-304
66. Zee DS. Perspectives on the pharmacotherapy of vertigo. Arch Otolaryngol. 1985; 111: 609-612
67. Curthoys IS, Halmagyi GM. Vestibular compensation: a review of the oculomotor, neural, and clinical consequences of unilateral vestibular loss. J Vestib Res 1995; 5: 67-107

68. Lacour M. Restoration of nerve function: Models and concepts. In: Lacour M, Toupet M, Denise P, Christen Y (Hrsg). *Vestibular compensation: Facts, theories and clinical perspectives*. Paris: Elsevier; 1989: 11-34
69. Curthoys IS, Halmagyi GM. Vestibular compensation. In: Büttner U (Hrsg). *Vestibular dysfunction and its therapy*. Adv Otorhinolaryngol 1999; 55: 82-110
70. Cawthorne T. Vestibular injuries. Proc R Soc Med 1946; 39: 270-273
71. Cooksey FS. Rehabilitation in vestibular injuries. Proc R Soc Med 1946; 39: 273-275
72. Sterkers JM. La méthode du point de mire pur la rééducation anti-vertigineuse. Rev Laryngol (Paris) 1977; 98: 535-539
73. Hamann KF. Training gegen Schwindel. Berlin: Springer, 1987
74. Hamann KF. Rehabilitation von Patienten mit vestibulären Störungen. HNO 1988; 36: 305-307
75. Gans RE. Vestibular Rehabilitation: Protocols and programs. San Diego: Singular Publishing Group, 1996
76. Cesarini A, Alpini D. Vertigo and dizziness rehabilitation. Berlin: Springer, 1999
77. Strupp M, Arbusow V, Brandt T. Exercise and drug therapy alter recovery from labyrinth lesion in humans. Ann N Y Acad Sci 2001; 942: 79-94
78. Strupp M, Arbusow V, Maag KP, Gall C, Brandt T. Vestibular exercises improve central vestibulospinal compensation after vestibular neuritis. Neurology 1998; 51: 838-844
79. Jerram AH, Darlington CL, Smith PF. Methylprednisolone reduces spontaneous nystagmus following unilateral labyrinthectomy in guinea pig. Eur J Pharmacol 1995; 275: 291-293
80. Pfaltz CR, Meran A. Sudden unilateral loss of vestibular function. Adv Oto Rhinol Laryng 1981; 27: 159-167
81. Pfaltz CR. Vestibular compensation. Acta Otolaryngol 1983; 95: 402-406
82. Stoll W. Die Begutachtung vestibulärer Störungen. Laryng Rhinol 1979; 58: 509-515
83. Stoll W. Untersuchungsmethoden zur Objektivierung und Begutachtung vestibulärer Störungen. Akt Neurol 1982; 9: 121-125
84. Stoll W. Das neurootologische Gutachten: Konsensusbericht und Fazit. In: Stoll W (Hrsg). *Das neurootologische Gutachten*. Stuttgart: Thieme, 2002; 146-155
85. Krückels G, Dujardin H, Westhofen M. Verfahren zur Abschätzung der vestibulären Kompensationsleistung-Klinische Beobachtung/quantitative Neurootometrie? Laryngo Rhino Otol 1999; 78: 204-209
86. Kley W, Reinecken R. Drehpendelprüfung (Eine Drehschwachreizprüfmethode des Vestibularapparates). Z Laryngol Rhinol Otol 1961; 40: 68-77
87. Söllner C, Burghammer M, Busch-Neunwisch E, Berger J, Schwarz H, Riekel C, Nicolson T. Control of crystal size and lattice formation by starmaker in oto-lith biomineralization. Science. 2003; 302: 282-286
88. Kawamata S, Igarashi Y. Growth and turnover of rat otoconia as revealed by labeling with tetracycline. Anat Rec 1995; 242: 259-266
89. Ross MD, Johnsson LG, Peacor D, Allard LF. Observation on normal and degenerating human otoconia. Ann Otol 1976; 85: 310-326
90. Lim DJ. Formation and fate of the otoconia. Ann Otol 1973; 82, 23-35
91. Takumida M, Zhang DM, Yajin K. In vitro calcium ion turnover in otoconia of the guinea pig. Auris Nasus Larynx 1997; 24: 119-124
92. Takumida M, Zhang DM, Yajin K, Harada Y. Polychromatic labeling of otoconia for the investigation of calcium turnover. ORL J Otorhinolaryngol Relat Spec 1997; 59: 4-9
93. Parnes LS, Agrawal SK, Atlas J. Diagnosis and management of benign paroxysmal positional vertigo (BPPV). CMAJ 2003; 30: 169: 681-693
94. Zucca G, Valli S, Valli P, Mira E (1998): Why do benign paroxysmal positional vertigo episodes recover spontaneously? J Vestibul Res 1998; 8: 325-329
95. Lindsay JR, Hemenway WG. Postural vertigo due to unilateral sudden partial loss of vestibular function. Ann Otol Rhinol Laryngol. 1956; 65: 692-706
96. Lindsay JR. Paroxysmal postural vertigo and vestibular neuronitis. Arch Otolaryngol. 1967; 85: 544-547
97. Parnes LS, McClure JA: Free floating endolymph particles: A new operative finding during posterior semicircular canal occlusion. Laryngoscope 1992; 102: 988-992
98. Welling DB, Parnes LS, O'Brien B, Bakalez LO, Brackmann DE, Hinojosa R. Particulate matter in the posterior semicircular canal. Laryngoscope 1997; 107:90-94
99. Walther LE. Benigner paroxysmaler Lagerungsschwindel. Forum HNO 2002; 5: 165-170
100. Schmäl F, Stoll W. Diagnostik und Therapie des benignen paroxysmalen Lagerungsschwindels Laryngo Rhino Otol 2002; 81: 368-380
101. von Brevern M, Lempert T. Benigner paroxysmaler Lagerungsschwindel. HNO 2002; 50: 671-681
102. Korres S, Balatsouras DG, Kaberos A, Economou C, Kandilatos D, Ferekidis E. Occurrence of semicircular canal involvement in benign paroxysmal positional vertigo. Otol Neurotol 2002; 23: 926-932
103. Bertholon P, Bronstein AM, Davies RA, Rudge P, Thilo KV. Positional down beating nystagmus in 50 patients: cerebellar disorders and possible anterior semicircular canalithiasis. J Neurol Neurosurg Psychiatry 2002; 72: 366-372
104. Steddin S, Brandt T. Unilateral mimicking bilateral benign paroxysmal positioning vertigo. Arch Otolaryngol Head Neck Surg. 1994; 120: 1339-1341
105. Sakata E, Uchida Y, Nakano Y, Takahashi K: Pathophysiology of positional vertigo of the malignant paroxysmal

- type. *Auris Nasus Larynx* 1984; 11: 79-90
106. Sakata E, Ohtsu K, Shimura H, Skai S. Positional nystagmus of benign paroxysmal type (BPPV) due to cerebellar vermis lesions: pseudo-BPPV. *Auris Nasus Larynx* 1987; 14: 17-21
107. Califano L, Montanaro SC, Capparuccia PG, Di Maria D, Villari G. Case report: ipsilateral association of posterior and anterior canalolithiasis *Acta Otorhinolaryngol Ital.* 2002; 22: 376-379
108. Steddin S, Brandt T. Benigner paroxysmaler Lagerungsschwindel. *Nervenarzt* 1994; 65: 505-510
109. Bertholon P, Faye MB, Tringali S, Martin Ch. Benign paroxysmal positional vertigo of the horizontal canal. Clinical features in 25 patients. *Ann Otolaryngol Chir Cervicofac.* 2002; 119: 73-80
110. Epley JM. Human experience with canalith repositioning maneuvers. *Ann N Y Acad Sci* 2001; 942: 179-191
111. Semont A, Freyss G, Vitte E: Curing the BPPV with a liberatory maneuver. *Adv Otorhinolaryngol* 1988; 42: 290-293
112. Epley JM: The canalith repositioning procedure: for treatment of benign paroxysmal positioning vertigo. *Otolaryngol Head Neck Surg* 1992; 107, 399-404
113. Hain TC, Helminski JO, Reis IL, Uddin MK. Vibration does not improve results of the canalith repositioning procedure. *Arch Otolaryngol Head Neck Surg* 2000; 126: 617-622
114. Li JC. Mastoid oscillation: a critical factor for success in canalith repositioning procedure. *Otolaryngol Head Neck Surg* 1995; 112: 670-675
115. Brandt T, Daroff RB. Physical therapy for benign paroxysmal positional vertigo. *Arch Otolaryngol.* 1980; 106: 484-485
116. Lempert T, Thiel-Wilck K: A positional maneuver for treatment of horizontal-canal benign positional vertigo. *Laryngoscope* 1996; 106: 476-478
117. Vannucchi P, Giannoni B, Pagnini P. Treatment of horizontal semicircular canal benign paroxysmal positional vertigo. *J Vestib Res* 1997; 7: 1-6
118. Cohen HS, Kimball KT. Treatment variations on the Epley maneuver for benign paroxysmal positional vertigo. *Am J Otolaryngol* 2004; 25: 33-37
119. Iro H, Wald Fahrer F, Wolf SR, Gjuric M, Haid CT, Wigand ME. Chirurgische Therapieoptionen bei Schwindel. In Stoll W (Hrsg). *Vestibuläre Erkrankungen-eine interdisziplinäre Herausforderung.* Thieme: Stuttgart, 2001; 77-82
120. Haid CT. Leitsymptome beim Akustikusneurinom. In: Stoll W. *Differenzialdiagnose Schwindel.* Springer Berlin, Heidelberg 1998; 29-37
121. Silverstein H. Surgery for vertigo (you don't have to live with it). *J Otolaryngol.* 1981; 10: 343-358
122. Gacek RR. Further observations on posterior ampullary nerve transection for positional vertigo. *Ann Otol Rhinol Laryngol* 1978; 87: 300-305
123. Bergström B. Morphology of the vestibular nerve. I. Anatomical studies of the vestibular nerve in man. *Acta Otolaryngol* 1973; 76: 162-172
124. Leuwer RM, Westhofen M. Surgical anatomy of the singular nerve. *Acta Otolaryngol* 1996; 116: 576-580
125. Nomura Y, Ooki S, Kukita N, Young YH. Laser labyrinthectomy. *Acta Otolaryngol* 1995; 115: 158-161
126. Gacek RR. Technique and results of singular neurectomy for the management of benign paroxysmal positional vertigo. *Acta Otolaryngol* 1995; 115: 154-157
127. Agrawal SK, Parnes LS. Human experience with canal plugging. *Ann N Y Acad Sci* 2001; 942: 300-305
128. Parnes LS, McClure JA. Posterior semicircular canal occlusion for intractable benign paroxysmal positional vertigo. *Ann Otol Rhinol Laryngol* 1990; 99: 330-334
129. Parnes LS, McClure JA. Posterior semicircular canal occlusion in the normal hearing ear. *Otolaryngol Head Neck Surg* 1991; 104: 52-57
130. Parnes LS. Update on posterior canal occlusion for benign paroxysmal positional vertigo. *Otolaryngol Clin North Am* 1996; 29: 333-342
131. Strutz J, Menke K. Die Chirurgie des benignen paroxysmalen Lagerungsschwindels. (Abstract) *HNO* 1998; 46: 417
132. Walsh RM, Bath AP, Cullen JR, Rutka JA. Long-term results of posterior semicircular canal occlusion for intractable benign paroxysmal positional vertigo. *Clin Otolaryngol* 1999; 24: 316-323
133. Brantberg K, Bergenius J. Treatment of anterior benign paroxysmal positional vertigo by canal plugging: a case report. *Acta Otolaryngol* 2002; 122: 28-30
134. Antonelli PJ, Lundy LB, Kartush JM, Burgio DL, Graham MD. Mechanical versus CO<sub>2</sub> laser occlusion of the posterior semicircular canal in humans. *Am J Otol* 1996; 17: 416-420
135. Wilpizeski C. Experimental labyrinthotomy in monkeys by argon and carbon dioxide lasers. *Otolaryngol Head Neck Surg.* 1981; 89: 197-203
136. Anthony PF. Partitioning of the labyrinth: application in benign paroxysmal positional vertigo. *Am J Otol.* 1991; 12: 388-593
137. Nomura Y. Argon laser irradiation of the semicircular canal in two patients with benign paroxysmal positional vertigo. *J Laryngol Otol* 2002; 116: 723-725
138. Anthony PF. Utricular macular ablation for benign paroxysmal positional vertigo. *Ear Nose Throat J* 1996; 75: 416-421
139. Basser LS. Benign paroxysmal vertigo of childhood (a variete of vestibular neuronitis). *Brain* 1964; 87: 141-152
140. Herraiz C, Calvin FJ, Tapia MC, de Lucas P, Arroyo R. The migraine: benign paroxysmal vertigo of childhood complex. *Int Tinnitus J* 1999; 5: 50-52
141. Drigo P, Carli G, Laverda AM. Benign paroxysmal vertigo of childhood. *Brain Dev* 2001; 23: 38-41
142. Uneri A, Turkdogan D. Evaluation of vestibular functions in children with

- vertigo attacks. *Arch Dis Child.* 2003; 88: 510-551
143. Hamann KF, von Czetylitz G. Neue Aspekte zur Differenzialdiagnose des kindlichen Schwindels. *HNO* 1987; 35: 267-269
144. Hilton M, Pinder D. Benign paroxysmal positional vertigo. *BMJ* 2003; 326: 673
145. Sakata E. Das neurootologische Studium über die Läsion des Kleinhirnwurms. *Equilibrium Res* 1971; Suppl 1: 30-48
146. Sakata E, Uchida Y, Nakano Y, Takahashi K. Pathophysiology of positional vertigo of the malignant paroxysmal type. *Auris Nasus Larynx* 1984; 11: 79-90
147. Sakata E, Ohtsu K, Shimura H, Skai S. Positional nystagmus of benign paroxysmal type (BPPV) due to cerebellar vermis lesions: pseudo-BPPV. *Auris Nasus Larynx* 1987; 14, 17-21
148. Del Rio M, Arriaga MA. Benign positional vertigo: prognostic factors. *Otolaryngol Head Neck Surg.* 2004; 130: 426-429
149. Hilton M, Pinder D. The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev* 2004; 2: CD003162
150. Asawachianginda S, Isipradit P, Snidvongs K, Supiyaphun P. Canalith repositioning for benign paroxysmal positional vertigo: a randomized, controlled trial. *Ear Nose Throat J* 2000; 79: 732-734, 736-737
151. Woodworth BA, Gillespie MB, Lambert PR. The canalith repositioning procedure for benign positional vertigo: a meta-analysis. *Laryngoscope* 2004; 114: 1143-1146
152. Okaja M, ebm 00858 (038.072) In: The Cochrane Library, 2004: 4. <http://www.ebm-guidelines.com>
153. Li JC, Li CJ, Epley J, Weinberg L. Cost-effective management of benign positional vertigo using canalith repositioning. *Otolaryngol Head Neck Surg* 2000; 122: 334-339
154. Stoll W. Morbus Menière. In: Stoll W (Hrsg). *Differenzialdiagnose Schwindel.* Berlin: Springer, 1998, 19-25
155. Michel O. *Morbus Menière.* Stuttgart: Thieme, 1997
156. Silverstein H, Hyman SM, Feldbaum J, Silverstein D. Use of streptomycin sulfate in the treatment of Menière's disease. *Otolaryngol Head Neck Surg* 1984; 92: 229-32
157. Schaaf H, Seling B, Rienhoff NK, Lambert A, Nelting M, Hesse G. Sind rezidivierende Tiefotonhörverluste -ohne Schwindel- die Vorstufe eines Morbus Menière? *HNO* 2001; 49: 543-547
158. Rask-Andersen H, Stahle J. Immuno-defence of the inner ear? Lymphocyte-macrophage interaction in the endolymphatic sac. *Acta Otolaryngol* 1980; 89: 283-294
159. Arnold W, Altermatt HJ, Gebbers JO. Qualitativer Nachweis von Immunoglobulinen im menschlichen Saccus endolymphaticus. *Laryngo Rhino Otol* 1984; 63: 464-467
160. Paparella MM, Djalilian HR. Etiology, pathophysiology of symptoms, and pathogenesis of Menière's disease. *Otolaryngol Clin North Am* 2002; 35: 529-545
161. Barbara M, Attanasio G, Petrozza V, Modesti A, Filipo R. The endolymphatic sac as the immunocompetent organ of the inner ear. *Ann N Y Acad Sci* 1997; 830: 243-252
162. Angelborg C. Pharmacological treatment of Meniere's disease. A review of common remedies with special emphasis on vasodilators, including hyperosmolar substances and hearing. *Scand Audiol Suppl* 1986; 26: 33-36
163. Arenberg IK, Lemke C, Shambaugh GE Jr. Viral theory for Meniere's disease and endolymphatic hydrops: overview and new therapeutic options for viral labyrinthitis. *Ann N Y Acad Sci.* 1997; 830: 306-313
164. Arnold W, Niedermeyer HP. Herpes simplex virus antibodies in the perilymph of patients with Menière disease. *Arch Otolaryngol Head Neck Surg* 1997; 123: 53-56
165. Morrison AW. Anticipation in Meniere's disease. *J Laryngol Otol* 1995; 109: 499-502
166. Düwel P, Jüngling E, Westhofen M, Lückhoff A. Potassium currents in vestibular type II hair cells activated by hydrostatic pressure. *Neuroscience* 2003; 116: 963-972
167. Arab SF, Düwel P, Jüngling E, Westhofen M, Lückhoff A. Inhibition of voltage-gated calcium currents in type II ves-tibular hair cells by cinnarizine. *Naunyn Schmiedebergs Arch Pharmacol* 2004; 369: 570-575
168. Haid CT. Evaluation of flunarizine in patients with Meniere's disease. Subjective and vestibular findings. *Acta Otolaryngol Suppl* 1988; 460: 149-153
169. Jäger L, Strupp M, Brandt T, Reiser M. Bildgebung von Labyrinth und Nervus vestibularis. *Nervenarzt* 1997; 68: 443-458
170. Krombach GA, Schmitz-Rode T, Tacke J, Düwel P, DiMartino E, Engelke JC, Günther RW, Westhofen M. Kernspintomografische Darstellung des Innenohrs bei Patienten mit sensorineuralem Hörverlust und Schwindel. *Laryngo Rhino Otol* 2001; 80: 177-181
171. Fujita N, Nario K, Yamanaka T, Ueda T, Matsunaga T. Neurovascular compression syndrome with Meniere's disease - a report of the treatment of a patient with microvascular decompression. *Auris Nasus Larynx* 1998; 25: 203-207
172. Committee on Hearing and Equilibrium of American Academy of Otolaryngology. Guidelines for diagnosis and evaluation for the therapy in Meniere's disease. *Otolaryngol Head Neck Surg* 1995; 113: 1881-1885
173. Waldfahrer F, Finke C, Iro H. Aktuelle Aspekte der medikamentösen Schwindeltherapie. In: Stoll W (Hrsg). *Vestibuläre Erkrankungen.* Stuttgart: Thieme, 2001: 98-107
174. Schmäl F, Stoll W. Medikamentöse Schwindeltherapie. *Laryngo Rhino Otolologie* 2003; 28: 44-66

175. Deering RB, Prescott P, Simmons RL, Downey LJ. A double-blind crossover study comparing betahistine and cinnarizine in the treatment of recurrent vertigo in patients in general practice. *Curr Med Res Opin* 1986; 10: 209-214
176. Novotny M, Kostrica R. Fixed combination of cinnarizine and dimenhydrinate versus betahistine dimesylate in the treatment of Meniere's disease: a randomized, double-blind, parallel group clinical study. *Int Tinnitus J* 2002; 8: 115-123
177. Laurikainen E, Miller JM, Nuttall AL, Quirk WS. The vascular mechanism of action of betahistine in the inner ear of the guinea pig. *Eur Arch Otorhinolaryngol* 1998; 255: 119-123
178. Kawabata A, Sasa M, Kishimoto T, Takaori S. Effects of anti-vertigo drugs on medial vestibular nucleus neurons activated by horizontal rotation. *Jpn J Pharmacol* 1991; 55: 101-106
179. James AL, Burton MJ. Betahistine for Meniere's disease or syndrome. *Cochrane Database Syst Rev* 2001; 1: CD001873
180. Mierzwiński J, Kazmierczak H, Pawlak-Osinska K, Piziewicz A. The effect of betahistine on vestibular habituation: comparison of rotatory and sway habituation training. *Acta Otolaryngol* 2001; 121: 610-615
181. van Deelen GW, Huizing EH. Use of a diuretic (Dyazide) in the treatment of Meniere's disease. A double-blind cross-over placebo-controlled study.
182. ORL J Otorhinolaryngol Relat Spec 1986; 48: 287-292
183. van de Heyning PH. Therapeutic strategies for vertigo in Meniere's disease. Abstract 2nd vertigo day Berlin, 1998
184. Vollrath M, Marangos N, Hesse G. Die Dehydratationstherapie des Tiefoton-hörverlustes. *HNO* 1990; 38:154-157
185. Pappas DG, Banyas JB. A newly recognized etiology of Meniere's syndrome. A preliminary report. *Acta Otolaryngol Suppl* 1991; 485: 104-107
186. Vischer M, Arnold W. Kortisonsensible Innenohrscherhörigkeit. *Otorhinolaryngol Nova* 1991; 1: 75-79
187. Niedermeyer HP, Zahneisen G, Lappa P, Busch R, Arnold W. Cortisol levels in the human perilymph after intravenous administration of prednisolone. *Audiol Neurotol* 2003, 8: 316-321
188. Waksman SA, Bugie E, Schatz A. Isolation of antibiotic substances from soil microorganisms with spezial reference to streptothricin and streptomycin. *Proc Staff Meet Mayo Clin* 1944; 19: 537
189. Hinshaw HC, Feldman WH. Streptomycin in treatment of clinical tuberculosis: A preliminary report. *Proc Staff Meet Mayo Clin* 1945; 20: 313
190. Causse R, Gondet I, Vallancien B. Action vestibulaire de la streptomycine chez souris. *C Soc Biol (Paris)* 1948; 142: 747
191. Fowles EP. Streptomycin treatment of vertigo. *Trans Am Acad Ophthalmol Otolaryngol Head Neck Surg* 1948; 113: 181-185
192. Schuknecht HF. Ablation therapy for the relief of Meniere's disease. *Trans Am Laryngol Rhinol Otol Soc. (60th Meeting)* 1956; 589-600
193. Schuknecht HF. Ablation therapy in the management of Meniere's disease. *Acta Otolaryngol.* 1957; 47(Suppl 132):1-42
194. Weinstein MJ, Leudemann GM, Oden EM et al. Gentamicin: A new broad spectrum antibiotic complex. *Antimicrob Agents Chemother* 1963; 3: 1
195. Bagger-Sjöback D. Effect of streptomycin and gentamicin on the inner ear. *Ann NY Acad Sci* 1997; 830: 120-129
196. Holz E, Stange G, Terajama Y. Minderung der Ototoxizität basischer Streptomycetesantibiotika anhand tierexperimenteller Untersuchungen. Vortrag IV. Int Kongr F Infektionskrankheiten. München: Schattauer, 1966
197. Lange G. Isolierte medikamentöse Ausschaltung eines Gleichgewichtsorgans beim Morbus Menière mit Streptomycin-Ozothin. *Arch Klin Exp Ohr Nas u Kehlk Heilk* 1968; 191: 545
198. Lange G. Ototoxische Antibiotika in der Behandlung des Morbus Menière. *Therapiewoche* 1976; 26: 3366-3369
199. Schmidt CL, Beck C. Behandlung des Morbus Menière mit intratympanal appliziertem Gentamicin-Sulfat. *Laryng Rhinol Otol* 1980; 59: 804-807
200. Nedzelski JM, Schessel DA, Bryce GE, Pfleiderer AG. Chemical labyrinthectomy: local application of gentamicin for the treatment of unilateral Meniere's disease. *Am J Otol* 1992; 13: 18-22
201. Monsell EM, Shelton C. Labyrinthotomy with streptomycin infusion: early results of a multicenter study. *The LSI Multicenter Study Group. Am J Otol* 1992; 13: 416-422
202. Nedzelski JM. Intratympanic Gentamicin results of an ongoing prospective outpatient treatment regimen (AAO-HNS criteria). XV. World Congress of ORL, Istanbul 1993
203. Katzke D. Die intratympanale Gentamicinbehandlung bei Morbus Menière. *Laryng Rhinol Otol* 1982; 61: 4-8
204. Küppers P, Ahrens H, Blessing R. Die kontinuierliche intratympanale Gentamicin-Infusion beim Morbus Menière. *HNO* 1994; 42: 429-433
205. Küppers P, Bach-Quang M, Blessing R. Die Gentamicin-Titration mittels Infusionspumpe: Eine neue Form der Menière-Therapie. *HNO-Information* 1991; 16: 20
206. Jahnke K. Transtympanic application of Gentamicin with cochlea protection. In: Nudol JB (Hrsg): Second International Symposium on Meniere's disease. Kugler und Ghedini, Amsterdam, Berkeley, Milano 1989
207. Jahnke K, Arweiler D. Die intravenöse Gentamicintherapie bei beidseitigem M. Menière. *Laryngo Rhino Otol* 1997; 76: 519-522
208. Marks S, Arenberg IK, Hoffer ME. Round-Window-Mikrokatheter-assisti-

- erte Mikrodosierung von Gentamicin: Alternative in der Behandlung des Tinnitus bei Patienten mit Morbus Menière. *Laryngo Rhino Otol* 2000; 79: 327-331
209. Lange G. 27 Jahre Erfahrung mit der transtympanalen Aminoglykosid-Behandlung des Morbus Menière. *Laryngo Rhino Otol* 1995; 74: 720-723
210. Lange G. Die Gentamicin-Injektionstechnik. *HNO* 1998; 46: 1000-1002
211. Lange G. Transtympanic treatment for Menière's disease with Gentamicin-sulfate. In: Vosteen KH, Schuknecht H et al. (Hrsg). Menière's disease. Stuttgart: Thieme 1981
212. Schacht J, Weiner N. Aminoglycoside-induced hearing loss: a molecular hypothesis. *ORL J Otorhinolaryngol Relat Spec* 1986; 48: 116-123
213. Schacht J. Molecular mechanisms of drug-induced hearing loss. *Hear Res* 1986; 22: 297-304
214. Schacht J. Biochemical basis of Aminoglykoside ototoxicity. *Otolaryngol Clin North Am* 1993; 26: 845-856
215. Park J, Cohen G. Vestibular ototoxicity in check: effects of streptomycin on equilibrium and on ampullary dark cells. *Am J Otolaryngol* 1982; 6: 117-127
216. Zenner HP, Schacht J. Hearing loss caused by Aminoglykoside antibiotics: affect on the membrane component PIP2 in outer hair cells as the mechanism of action. *HNO*. 1986; 34: 417-423
217. Chia SH, Gamst AC, Anderson JP, Harris JP. Intratympanic Gentamicin Therapy for Meniere's disease: A meta-analysis. *Otol Neurotol* 2004; 25: 544-552
218. Pfeiderer AG. The current role of local intratympanic gentamicin therapy in the management of unilateral Meniere's disease. *Clin Otolaryngol* 1998; 23: 34-41
219. Atlas JT, Parnes LS. Intratympanic gentamicin titration therapy for intractable Meniere's disease. *Am J Otol* 1999; 20: 357-363
220. Martin E, Perez N. Hearing loss after intratympanic gentamicin therapy for unilateral Meniere's Disease. *Otol Neurotol* 2003; 24: 800-806
221. Quaranta A, Aloisi A, De Benedittis G, Scaringi A. Intratympanic therapy for Meniere's disease. High-concentration gentamicin with round-window protection. *Ann NY Acad Sci* 1999; 28: 884: 410-424
222. Thomsen J, Charabi S, Tos M. Preliminary results of a new delivery system for Gentamicin to the inner ear in patients with Menière's disease. *Eur Arch Otorhinolaryngol* 2000; 257: 362-365
223. Hoffer ME, Kopke RD, Weisskopf P, Gottshall K, Allen K, Wester D, Balaban C. Use of the round window microcatheter in the treatment of Meniere's disease. *Laryngoscope*. 2001; 111: 2046-2049
224. Minor LB. Intratympanic gentamicin for control of vertigo in Meniere's disease: vestibular signs that specify completion of therapy. *Am J Otol* 1999; 20: 209-219
225. Magnusson M, Padoan S, Karlberg M, Johansson R. Delayed onset of ototoxic effects of gentamicin in patients with Meniere's disease. *Acta Otolaryngol Suppl* 1991; 485: 120-122
226. Ödkvist LM, Bergenius J, Möller C. When and how to use gentamicin in the treatment of Meniere's disease. *Acta Otolaryngol Suppl.* 1997; 526: 54-57
227. Lange G, Mann W, Maurer J. Intratympanale Intervalltherapie des Morbus Menière mit Gentamicin unter Erhalt der Cochleafunktion. *HNO* 2003; 51: 598-902
228. Lange G, Maurer J, Mann W. Long-term results after interval therapy with intratympanic gentamicin for Meniere's disease. *Laryngoscope* 2004; 114:102-105
229. Hoffmann F, Beck C, Beck C, Stratulat S. Subablative Gentamicintherapie beim Morbus Menière. *HNO* 1991; 41: 296-300
230. Driscoll CL, Kasperbauer JL, Facer GW, Harner SG, Beatty CW. Low-dose intratympanic gentamicin and the treatment of Meniere's disease: preliminary results. *Laryngoscope*. 1997; 107: 83-89
231. Blakley BW. Update on intratympanic gentamicin for Meniere's disease. *Laryngoscope*. 2000; 110: 236-240
232. Shea JJ. Perfusion of the inner ear with streptomycin. *Am J Otol* 1989; 102: 150-155
233. Shea JJ, Norris CH. Streptomycin perfusion of the labyrinth. In: Proceedings of the second International Symposium on Menière's disease. Nadol JB (Hrsg). Amsterdam: Kugler & Ghedini, 1989
234. Wang D, Wang Z. Regeneration and functional restoration of vestibular hair cells in guinea pigs after gentamicin damage. *Zhonghua Er Bi Yan Hou Ke Za Zhi* 1997; 32: 153-156
235. Forge A, Li L, Nevill G. Hair cell recovery in the vestibular sensory epithelia of mature guinea pigs. *J Comp Neurol* 1998; 397: 69-88
236. De Waele C, Meguenni R, Freyss G, Zamith F, Bellalimat N, Vidal PP, Tran Ba Huy P. Intratympanic gentamicin injections for Menière disease: vestibular hair cell impairment and regeneration. *Neurology* 2002; 59: 1442-1444
237. Silverstein H, Rowan PT, Olds MJ, Rosenberg SI. Inner ear perfusion and the role of round window patency. *Am J Otol* 1997; 18: 586-589
238. Diamond C, O'Connell DA, Hornig JD, Liu R. Systematic review of intratympanic gentamicin in Menière's disease. *J Otolaryngol* 2003; 32: 351-361
239. Bárány R. Die Beeinflussung des Ohrensausens durch intravenös injizierte Localanästhetica. *Acta Otolaryngol* 1935; 23: 210-207
240. Gstöttner W, Adunka O. Labyrinthanästhesie bei M. Menière. *HNO* 2004; 52: 265
241. Ristow W. Zur Behandlung der Menière-Krankheit mittels temporärer Labyrinthanästhesie. *Z Laryngol Rhinol* 1968; 47: 442-448

242. Fradis M, Podoshin L, Ben-David J, Reiner B. Treatment of Menière's disease by intratympanic injection with lidocaine. *Arch Otolaryngol* 1985; 111: 491-493
243. Adunka O, Moustaklis E, Weber A, May A, von Ilberg C, Gstötzner W, Kierner AC. Labyrinth anesthesia - a forgotten but practical treatment option in Meniere's disease. *ORL J Otorhinolaryngol Relat Spec* 2003; 65: 84-90
244. Sakata E, Kitago Y, Murata Y, Teramoto K. Behandlung der Menière'schen Krankheit. Paukenhöhleinfusion von Lidocain- und Steroidlösung. *Auris Nasus Larynx* 1986; 13: 79-89
245. Plontke SK, Salt AN. Quantitative interpretation of corticosteroid pharmacokinetics in inner fluids using computer simulations. *Hear Res* 2003; 182: 34-42
246. Bachmann G, Su J, Zumege C, Wittekindt C, Michel O. Permeabilität der runden Fenstermembran für Prednisolon-21-Hydrogensuccinat. *HNO* 2001; 49: 538-542
247. Silverstein H. Use of a new device, the MicroWick® to deliver medication to the inner ear. *Ear Nose Throat J* 1999; 78: 595-598
248. Charabi S, Thomsen J, Tos M. Round window gentamicin mu-catheter-a new therapeutic tool in Menière's disease. *Acta Otolaryngol Suppl* 2000; 543: 108-110
249. Schwab B, Lenarz T, Heermann R. Der Round-Window μCath zur Lokaltherapie des Innenohres-Ergebnisse einer plazebokontrollierten, prospektiven Studie beim chronischen Tinnitus. *Laryngo Rhino Otol* 2004; 83: 164-172
250. Plontke SK, Plinkert PK, Plinkert B, Koitschev A, Zenner HP, Löwenheim H. Transtympanic endoscopy for drug delivery to the inner ear using a new microendoscope. *Adv Otorhinolaryngol* 2002; 59:149-155.
251. Goycoolea MV, Lundman L. Round window membrane. Structure function and permeability: a review. *Microsc Res Tech* 1997; 36: 201-211
252. Rarey KE, Curtis LM. Receptors for glucocorticoids in the human inner ear. *Otolaryngol Head Neck Surg* 1996; 115: 38-41
253. Hughes GB, Kinney SE, Barna BP, Calabrese LH. Practical versus theoretical management of autoimmune inner ear disease. *Laryngoscope* 1984; 94: 758-767
254. McCabe BF. Autoimmune sensorineural hearing loss. 1979. *Ann Otol Rhinol Laryngol* 2004; 113: 526-530
255. Hillman TM, Arriaga MA, Chen DA. Intratympanic steroids: do they acutely improve hearing in cases of cochlear hydrops? *Laryngoscope*. 2003; 113: 1903-1907
256. Parnes LS, Sun AH, Freeman DJ. Corticosteroid pharmacokinetics in the inner ear fluids: an animal study followed by clinical application. *Laryngoscope* 1999; 109: 1-17
257. Dodson KM, Woodson E, Sismanis A. Intratympanic steroid perfusion for the treatment of Meniere's disease: a retrospective study. *Ear Nose Throat J* 2004; 83: 394-398
258. Barrs DM, Keyser JS, Stallworth C, McElveen JT Jr. Intratympanic steroid injections for intractable Meniere's disease. *Laryngoscope* 2001; 111: 2100-2104
259. Plontke SK, Salt AN. Quantitative interpretation of corticosteroid pharmacokinetics in inner fluids using computer simulations. *Hear Res* 2003; 182: 34-42
260. Alzamil KS, Linthicum FH Jr. Extraneous round window membranes and plugs: possible effect on intratympanic therapy. *Ann Otol Rhinol Laryngol* 2000; 109: 30-32
261. Arslan M. Erwägungen über die Wahl des chirurgischen Verfahrens bei der Menière'schen Krankheit. Die neue Methode der "osmotischen Induktion". *HNO* 1971; 19: 81- 87
262. Hüttenbrink KB. Vestibuläre Symptome durch mechanische Interaktionen zwischen Mittel- und Innenohr. In: Stoll W (Hrsg). Differenzialdiagnose Schwindel. Berlin: Springer, 1998: 85-90
263. Hüttenbrink KB. Die Bewegungen von Stapes-Piston-Prothesen bei Änderungen des statischen Luftdruckes. *Laryng Rhinol Otol* 1988; 67: 250-244
264. Hüttenbrink KB. Die Mechanik der Gehörknöchelchen bei statischen Drücken. II. Behinderte Gelenkfunktion und operative Kettenrekonstruktion. *Laryng Rhinol Otol* 1988; 67: 100-105
265. Hüttenbrink KB. Die Funktion der Gehörknöchelchenkette und der Muskeln des Mittelohres. *Eur Arch Oto Rhino Laryngol Suppl* 1995; 1-52
266. Tumarkin A. Thoughts on the treatment of labyrinthopathy. *J Laryngol Otol* 1966; 80: 1041-1053
267. Lall M. Meniere's disease and the grommet (a survey of its therapeutic effects). *J Laryngol Otol* 1969; 83: 787-791
268. Thomsen J, Bonding P, Becker B, Stage J, Tos M. The non-specific effect of endolymphatic sac surgery in treatment of Meniere's disease: a prospective, randomized controlled study comparing „classic“ endolymphatic sac surgery with the insertion of a ventilating tube in the tympanic membrane. *Acta Otolaryngol* 1998; 118: 769-773
269. Montandon P, Guillemin P, Häusler R. Prevention of vertigo in Meniere's syndrome by means of transtympanic ventilation tubes. *ORL J Otorhinolaryngol Relat Spec* 1988; 50: 377-381
270. Häusler R, Guillemin P, Montandon P. Surgical treatment of Meniere's disease by sacculotomy, cochleo-sacculotomy and transtympanic aerators. *Rev Laryngol Otol Rhinol (Bord)*; 1991: 112: 149-152
271. Hall MC, Brackmann DE. Eustachian tube blockage and Meniere's disease. *Arch Otolaryngol* 1977; 103: 355-357
272. Franz P, Hamzavi JS, Schneider B, Ehrenberger K. Do middle ear muscles trigger attacks of Meniere's disease? *Acta Otolaryngol* 2003 123: 133-137

273. Kitahara M, Kodama A, Ozawa H, Izukura H. Pressure test for the diagnosis of Meniere's disease. *Acta Otolaryngol Suppl* 1994; 510: 107-110
274. Konradsson KS, Svensson O, Carlborg B, Grenner J. Tympanic pressure gradients and otoacoustic emissions. *Ear Hear* 1999; 20: 403-409
275. Konradsson KS, Carlborg B, Grenner J, Tjernström O. Electrocochleographic and audiometric evaluation of hypobaric effect in Meniere's disease. *Laryngoscope* 1999; 109: 59-64
276. Ödkvist LM, Arlinger S, Billermark E, Densert B, Lindholm S, Wallqvist J. Effects of middle ear pressure changes on clinical symptoms in patients with Meniere's disease--a clinical multicentre placebo-controlled study. *Acta Otolaryngol Suppl* 2000; 543: 99-101
277. Densert B, Densert O, Arlinger S, Sass K, Ödkvist L. Immediate effects of middle ear pressure changes on the electrocochleographic recordings in patients with Meniere's disease: a clinical placebo-controlled study. *Am J Otol* 1997; 18: 726-733
278. Densert B, Arlinger S, Ödkvist LM. New technology to control symptoms in Meniere's disease. *Acta Otolaryngol* 2000; 120: 672-674
279. Densert B, Sass K. Control of symptoms in patients with Meniere's disease using middle ear pressure applications: two years follow-up. *Acta Otolaryngol* 2001; 121: 616-621
280. van Deelen GW, Hulk J, Huizing EH. The use of the underpressure chamber in the treatment of patients with Meniere's disease. *J Laryngol Otol*. 1987; 101: 229-35
281. Gruber J. Über Morbus Menière. *Mschr Ohrenheilk* 1895; 29: 181-184
282. Lacher G. Current historical perspectives on endolymphatic sac surgery. *Acta Otolaryngol Stockh Suppl* 1997; 526: 50-53
283. Portmann G. Recherches sur la physiologie du sac et du canal endolymphatiques: valeur fonctionnelle de l'organe endolymphatique des Sélaciens. *C R Soc Biol* 1921; 85: 1070-1077
284. Portmann G. Le traitement chirurgical des vertiges par l'ouverture du sac endolymphatique. 1926, Paris: Presse Médical
285. Portmann G. The saccus endolymphaticus and an operation for draining the same for the relief of vertigo. *Laryngol* 1927; 42: 809-817
286. Paparella MM, Sajjadi H. Endolymphatic sac enhancement. Principles of diagnosis and treatment. *Am J Otol* 1987 8: 294-300
287. Paparella MM, Sajjadi H. Endolymphatic sac revision for recurrent Meniere's disease. *Am J Otol*. 1988 9: 441-447
288. Paparella MM, Hamid RD. Etiology, pathophysiology of symptoms, and pathogenesis of Menière's disease. *Otolaryngol Clin N Am* 2002; 35: 529-545
289. Paparella MM. Revision of endolymphatic sac surgery for recurrent Menieres disease. *Otolaryngol Clin N Am* 2002; 35: 607-619
290. House WF. Subarachnoid shunt for drainage of endolymphatic hydrops. A preliminary report. *Laryngoscope* 1962; 72: 713-729
291. Plester D. Surgical treatment of Meniere's disease. *HNO (Berl)* 18: 205-210
292. Strutz J, Mann W (Hrsg). *Praxis der HNO-Heilkunde, Kopf- und Halschirurgie*. Stuttgart: Thieme, 2000
293. Thomsen J, Bretlau P, Tos M, Johnsen NJ. Placebo effect in surgery for Menière's disease. *Arch Otolaryngol* 1981; 107: 271-277
294. Thomsen J, Bretlau P, Tos M, Johnsen NJ. Menière's disease: a 3-year follow-up of patients in a double-blind placebo-controlled study on endolymphatic sac shunt surgery. *Adv Oto Rhino Laryng* 1983; 30: 350-354
295. Thomsen J, Bretlau P, Tos M, Johnsen NJ. Endolymphatic sac suc-mastoid shunt surgery. A non-specific treatment modality? *Ann Otol Rhinol Laryngol* 1986; 95: 32-35
296. Bretlau P, Thomsen J, Tos M, Johnsen NJ. Placebo effect in surgery for Menière's disease: nine-year follow-up. *Am J Otol* 1989; 10: 259-261
297. Thomsen J, Kerr A, Bretlau P, Olsson J, Tos M. Endolymphatic sac surgery: why we do not do it. The non-specific effect of sac surgery. *Clin Otolaryngol*. 1996; 21: 208-211
298. Vaisrub N. Summary statement to "placebo effect insurgery for Menière's disease sac shunt surgery disputed". (Letter to the editor). *Arch Otolaryngol* 1981; 107: 174
299. Pillsbury HC 3rd, Arenberg IK, Ferraro J, Ackley RS. Endolymphatic sac surgery. The Danish sham surgery study: an alternative analysis. *Otolaryngol Clin North Am* 1983 16: 123-127
300. Huang TS, Lin CC, Chang YL. Endolymphatic sac surgery for Meniere's disease. A cumulative study of twelve years' experience. *Acta Otolaryngol Suppl.* 1991; 485: 145-154
301. Kerr AG, Toner JG, McKee GJ, Smyth GD. Role and results of cortical mastoidectomy and endolymphatic sac surgery in Meniere's disease. *J Laryngol Otol* 1989; 103: 1161-1166
302. Silverstein H, Smouha E, Jones R. Natural history vs. surgery for Menière's disease. *Otolaryngol Head Neck Surg* 1989; 100: 6-16
303. Schuknecht HF, Rüther A. Blockage of longitudinal flow in endolymphatic hydrops. *Eur Arch Otolaryngol* 1990; 248: 209-217
304. Wilschowitz M, Sanchez-Hanke M, Ußmüller J. Zur Wertigkeit der Saccotomie beim M. Menière – Eine Langzeitanalyse von 42 Fällen. *HNO* 2001; 49: 180-187
305. Ußmüller J, Sanchez-Hanke R, Leuwer R. Klinische Relevanz entwicklungsbedingter Veränderungen des Saccus endolymphaticus. *Laryngo Rhino Otol* 2001; 80: 308-312
306. Welling DB, Nagaraja HN. Endolymphatic mastoid shunt: a reevaluation

- of efficacy. *Otolaryngol Head Neck Surg* 2000; 122: 340-345
307. Pensak ML, Friedman RA. The role of endolymphatic mastoid shunt surgery in the managed care era. *Am J Otol*. 1998; 19: 337-340
308. Kato BM, LaRouere MJ, Bojrab DI, Michaelides EM. Evaluating quality of life after endolymphatic sac surgery: The menière's disease outcomes questionnaire. *Otol Neurotol* 2004; 25: 339-344
309. Huang TS. Endolymphatic sac surgery for Meniere's disease: experience with over 3000 cases. *Otolaryngol Clin North Am* 2002; 35: 591-606
310. Stahle J, Friberg U, Svedberg A. Long-term progression of Menierè's disease. *Acta Otolaryngol (Stockh) Suppl* 1991; 485: 78-83
311. Huang TS, Lin CC. Endolymphatic sac ballooning surgery for Meniere's disease. *Ann Otol Rhinol Laryngol* 1994; 103: 389-394
312. Huang TS, Hsu JC, Lee FP. Electrocochleographic monitoring in endolymphatic sac surgery for Menierè's disease. *Arch Otolaryngol Head Neck Surg* 1994; 120: 552-229
313. Bagger-Sjöback D. Surgical anatomy of the endolymphatic sac. *Am J Otol*. 1993; 14: 576-579
314. Yin S, Ke G, Gu N, Shen P, Zhang S, Wu Y, Zang W. Long-term results of surgical treatment of intractable Meniere's disease for control of vertigo. *Lin Chuang Er Bi Yan Hou Ke Za Zhi* 1999; 13: 291-292
315. Kitahara T, Horii A, Mishiro Y, Fukushima M, Kondoh K, Okumura S, Takeda N, Kubo T. Changes in plasma inner ear hormones after endolymphatic sac drainage and steroid-instillation surgery (EDSS) *Nippon Jibiinkoka Gakkai Kaiho* 2002; 105: 557-563
316. Huang TS. Topical mitomycin C and cephalosporin in endolymphatic sac surgery. *Laryngoscope* 2002; 112: 243-247
317. Welling DB, Pasha R, Roth LJ, Barin K. The effect of endolymphatic sac excision in Meniere disease. *Am J Otol* 1996; 17: 278-282
318. Westhofen M, Koch U. Therapieerfolge nach Saccotomie - Postoperativer Verlauf der vestibulären Kompensation. *HNO* 1988; 36: 315-317
319. Schwager K, Baier G, El-Din N, Shehata-Dieler W, Carducci F, Helms J. Revision surgery after saccotomy for Meniere's disease: does it make sense? *Eur Arch Otorhinolaryngol* 2002; 259: 239-242
320. Iro H, Waldfahrer F. Chirurgische Therapie bei Morbus Menière. In: Haid CT (Hrsg). *Schwindel aus interdisziplinärer Sicht*. Stuttgart: Thieme, 2003: 217-224
321. Schuknecht HF. Cochleosacculotomy for Meniere's disease: theory, technique and results. *Laryngoscope* 1982; 92: 853-858
322. Montandon PB, Häusler RJ, Kimura RS. Treatment of endolymphatic hydrops with cochleosacculotomy. Clinical results and experimental findings. *Otolaryngol Head Neck Surg* 1985; 93: 615-21.
323. Schuknecht HF. The cochlear endolymphatic (CE) shunt: Update on results. Spring Meeting of the American Neurotology Society, New Orleans, April 8, 1983
324. Kangsanarak J, Navacharoen N, Prawatmuang W, Supamanee T. Surgery for intractable vertigo: Chiang Mai experience. *J Med Assoc Thai* 1995; 78: 289-297
325. Kinney WC, Nalepa N, Hughes GB, Kinney SE. Cochleosacculotomy for the treatment of Meniere's disease in the elderly patient. *Laryngoscope* 1995; 105: 934-937
326. Rosenberg SI. Vestibular surgery for Meniere's disease in the elderly: a review of techniques and indications. *Ear Nose Throat J* 1999; 78: 443-446
327. Huang TS, Lin CC. Delayed endolymphatic hydrops: study and review of clinical implications and surgical treatment. *Ear Nose Throat J* 2001; 80:76-8, 81-82
328. Dandy WE. Menierè's disease. Its diagnosis and a method of treatment. *Arch Surg* 1928; 16: 1127-1152
329. Dandy WE. Menierè's disease: diagnosis and treatment. Report of thirty cases. *Am J Surg* 1933; 20: 693-398
330. McKenzie KG. Intracranial division of the vestibular portion of the auditory nerve for Menierè's disease. *CMAJ* 1936; 34: 369-381
331. House WF. Surgical exposure of the internal auditory canal and its contents through the middle cranial fossa. *Laryngoscope* 1961; 71: 1363-1385
332. Fisch U. Neurectomy of the vestibular nerve: surgical technique: indications and results obtained in 70 cases. *Rev Laryngol Otol Rhinol (Bord)* 1969; 90: 661-672
333. Wigand ME, Haid CT, Berg M, Rettinger G. The enlarged transtemporal approach to cerebello-pontine angle; technique and indications. *Acta Otorhinolaryngol (Ital)* 1982; 2: 571
334. Silverstein H. Transmeatal labyrinthectomy with and without and cochleovestibular neurectomy. *Laryngoscope* 1976; 36: 1777-1791
335. Fisch U. Die chirurgische Behandlung des Morbus Menière. *Arch Otolaryngol* 1976; 212: 385-391
336. Molony TB. Decision making in vestibular neurectomy. *Am J Otol* 1996; 17: 421-424
337. Thomsen JC, Berner AS, Tos M. Vestibular neurectomy in the treatment of Meniere's disease. *Ugeskr Laeger* 2000; 162: 5501-5503
338. Hillman TA, Chen DA, Arriaga MA. Vestibular nerve section versus intratympanic gentamicin for Meniere's disease. *Laryngoscope* 2004; 114: 216-22.
339. Rosenberg S, Silverstein H, Flanzer J, Wanamaker H. Bilateral Meniere's disease in surgical versus nonsurgical patients. *Am J Otol* 1991; 12: 336-340
340. Van de Heyning PH, Verlooy J, Schatteman I, Wyts FL. Selective vestibular neurektomy in Menierè's disease: a re-

- view. *Acta Otolaryngol (Stockh) Suppl* 1997; 526: 58-66
341. Silverstein H, Wanamaker H, Flanzer J, Rosenberg S. Vestibular neurectomy in the United States-1990. *Am J Otol* 1992; 13: 23-30
342. Schuknecht H. Ablation therapy für die relief of Menière's disease. *Laryngoscope* 1956; 66: 859-871
343. Cawthorne T. Membraneous labyrinthectomy via oval window for Menière's disease. *J Laryngol Otol* 1957; 71: 524-527
344. Helms J. Chirurgie an Labyrinth und innerem Gehörgang bei Gleichgewichtsstörungen. In: Naumann HH, Helms J, Herberhold C, Jahrsdoerfer RA, Kastenbauer ER, Panje WR, Tardy ME Jr (Hrsg). Kopf- und Hals-Chirurgie, Bd 2. Stuttgart: Thieme, 1996: 277-300
345. Jones R, Silverstein H, Smouha E. Long-term results of transmeatal cochleovestibular neurectomy: an analysis of 100 cases. *Otolaryngol Head Neck Surg* 1989; 100: 22-29
346. Peacock R. Alcoholic labyrinthine injection through the oval window in the treatment of aural vertigo. *Lancet* 1938; 1: 421-423
347. Mollison WW. Surgical treatment of vertigo by opening the external semicircular canal and injecting alcohol. *Lancet* 1938; 1: 421-423
348. Day KM. Labyrinth surgery for Menière's disease. *Laryngoscope* 1943; 53: 617-630
349. van Fick IA. Symposium: management of Meniere's disease. VI. Sacculotomy for hydrops. *Laryngoscope* 1965 Oct; 75(10): 1539-1546
350. Cody DT. The tack operation. *Arch Otolaryngol* 1973; 97: 109-11
351. Pulec JL. The otic-perotic shunt. *Otolaryngol Clin North Am* 1968; 1: 643-648
352. Arslan M. Direkte Applikation des Ultraschalls auf das knöcherne Labyrinth zur Therapie der Labyrinthose (Morbus Menière). *HNO* 1953-1954; 4: 166-168.
353. Arslan M, Giacomelli F, Mozzo W. Experimental reproduction of Menière's disease II. Observations on the permeability of the round window membrane. *Arch Ital Otol Rinol Laringol Suppl* 1963; 46: 21-26
354. Arslan M. Origin of the nystagmus arising during and after ultrasonic destructive irradiation of the vestibular apparatus. *Acta Otolaryngol Suppl* 1963; 192: 121
355. Arslan M. Ultrasonic destruction of the vestibular receptors in severe Menière's disease. *Laryngoscope*. 1964; 74: 1262-1282
356. Lermoyez M. Le vertige qui fait entendre (angiospasme labyrinthique). *Ann mal oreille Paris* 1929; 48: 575
357. Kempf HG, Jahnke K. Lermoyez-Syndrom. *HNO* 1989; 37: 276-280
358. Schmidt PH, Schoonhoven R. Lermoyez's syndrome. A follow-up study in 12 patients. *Acta Otolaryngol* 1989; 107: 467-473
359. Young YH, Wu CH. Electronystagmographic findings in a case of Lermoyez's syndrome. *Auris Nasus Larynx*. 1994; 21: 118-121
360. Maier W, Marangos N, Schendorff A. Das Lermoyez-Syndrom-Elektroneustagmographische Beobachtungen. *Laryngo Rhino Otol* 1996; 75: 372-376
361. Boenninghaus HG, Feldmann H, Steinmann G. Ein Beitrag zur Entstehung von Schwindel und Hörstörung bei Menière'scher Erkrankung und Lermoyezschem Syndrom. *Laryngol Rhinol Otol* 1967; 46: 709-727
362. Pillsbury HC 3rd, Postma DS. Lermoyez's syndrome and the otolithic crisis of Tumarkin. *Otolaryngol Clin North Am* 1983; 16: 197-203
363. Tumarkin A. The otolithic catastrophe. A new syndrome. *Br Med J* 1936; 175: 177
364. Huy PT, Toupet M. Peripheral disorders in the otolith system. A pathophysiological and clinical overview. *Adv Otorhinolaryngol* 2001; 58: 110-127
365. Ödkvist LM, Bergenius J. Drop attacks in Menière's disease. *Acta Otolaryngol (Stockh) Suppl* 1986; 455: 82-85
366. Westhofen M. Krankheitsbilder mit isolierten und kombinierten Otolithenfunktionsstörungen. In Haid CT. Schwindel aus interdisziplinärer Sicht. Stuttgart: Thieme, 2003, 153-162
367. Hofferberth B. Pathophysiologie und Klinik zentral-vestibulärer Erkrankungen. In: Haid CT (Hrsg). Schwindel aus interdisziplinärer Sicht. Stuttgart: Thieme, 2004: 199-205
368. Schuknecht HF. Delayed endolymphatic hydrops. *Ann Otol Rhinol Laryngol* 1978; 87: 743-748
369. Colebatch JG, Halmagyi GM, Skuse NF. Myogenic potentials generated by a click-evoked vestibulocollic reflex. *J Neurol Neurosurg Psychiatry* 1994; 57: 190-197
370. Westhofen M, Mangold U, Wöllmer W, Wehner M. Lasertherapie von Otolithenstörungen. In: Stoll W (Hrsg) Differenzialdiagnose Schwindel. Springer, Berlin 1998: 73-82
371. Nomura Y, Hara M, Okuno T. Application of argon laser to the inner ear. *Acta Otolaryngol*. 1988; 105: 439-444
372. Nomura Y, Okuno T, Young YH, Hara M. Laser labyrinthectomy in humans. *Acta Otolaryngol*. 1991; 111: 319-326
373. Nomura Y, Okuno T, Mizuno M. Treatment of vertigo using laser labyrinthectomy. *Acta Otolaryngol*. 1993; 113: 261-262
374. Clarke A. Sensory interaction in the vestibular system. Habilitationsarbeit für die Freie Universität Berlin 1992
375. Scherer H, Teiwes W, Clarke A. Measuring three dimensions of eye movement in dynamic situations by means of videooculography. *Acta Otolaryngol (Stockh)* 1991; 111: 182-187
376. Brandt T. Bilateral vestibulopathy revisited. *Eur J Med Res* 1995/96; 1: 361-368
377. Rinne T, Bronstein AM, Rudge P, Gresty MA, Luxon LM. Bilateral loss of vestibular function: clinical findings in 53 patients. *J Neurol* 1998; 245: 314-321

378. Bechterew W. Ergebnisse der Durchschneidung des Nervus acusticus, nebst Erörterung der Bedeutung der semicirculären Kanäle für das Körpergleichgewicht. *Pflügers Arch f d ges Physiologie* 1883; 30: 312-347
379. Schmidt CL, Maurer J. Bechterew-Phänomen beim Menschen. *Laryngo Rhino Otol* 1999; 78: 455-461
380. Dandy WE. The surgical treatment of Menière's disease. *Am J Surg* 1941; 20: 52
381. Lange G, Keller R. Beiderseitiger Funktionsverlust der peripheren Gleichgewichtsorgane. *Laryngo Rhino Otol* 2000; 79: 77-80
382. Reimann IR, Meier-Hellmann A, Pfeifer R, Traut T, Schilling A, Stein G, Reinhart K, Hoffmann A. Serumspiegelorientierte Dosierung der einmal täglichen Aminoglykosidtherapie beim kritisch Kranken: Ergebnisse einer prospektiven Untersuchung. *Anesthesiol Intensivmed Notfal* 1999; 34: 288-296
383. Orsoni JG, Zavota L, Pellistri I, Piazza F, Cimino L. Cogan syndrome. *Cornea*. 2002; 21: 356-359
384. Herdmann SJ. Vestibular rehabilitation. Philadelphia: Davis, 2000
385. Link W, Schneider D. Lebensqualität beim Altersschwindel in Deutschland. *Fortschritte der Medizin* 2003; 212: 47-50
386. Jeschke D, Zeilberger K. Altern und körperliche Aktivität. *Dtsch Arztebl* 2004; 101: 789-798
387. Judge JO, Whipple RH, Wolfson LI: Effects of resistive and balance exer-
- cises on isokinetic strength in older persons. *J Am Geriatr Soc* 1994; 42: 937-946
388. von Renteln-Kruse W, Micoll W, Oster P, Schlierf G. Arzneimittelverordnung, Schwindel und Stürze bei über 75jährigen Krankenhauspatienten. *Z Gerontol Geriat* 1998; 31: 286-289
389. Allum JHJ. Neue Ansätze zur Sturzprävention mittels Gleichgewichtsprothesen. Jahrestagung Deutsche Gesellschaft für Audiologie 2004; Leipzig, Vortrag
390. Noppens R, Hennes HJ. Kinetosen bei Hubschraubereinsätzen. *Notfall & Retlingsmedizin* 2001; 4: 426-430
391. Helling K, Hausmann S, Flöttmann T, Scherer H. Untersuchungen zur interindividuell unterschiedlichen Kinetoseempfindlichkeit. *HNO* 1997; 45: 210-215
392. Diamond SG, Markham CH. Validating the hypothesis of otolith asymmetry as a cause of space motion sickness. *Ann NY Acad Sci* 1992; 656: 725-731
393. Schmäl F, Stoll W. Kinetosen. *HNO* 2000; 48: 346-356